

On Convergence of Inverse Functions of Operators*

ARTHUR JAFFE, ANDRZEJ LESNIEWSKI, AND KONRAD OSTERWALDER†

Harvard University, Cambridge, Massachusetts 02138

Communicated by L. Gross

Received September 25, 1987

We study sequences $\{A_n\}$ of self-adjoint operators on a Hilbert space \mathcal{H} . We give a sufficient condition on a function f and on the $\{A_n\}$ such that $f(|A_n|) \rightarrow f(|A|)$ ensures $A_n \rightarrow A$. © 1988 Academic Press, Inc.

Let $\{A_n\}_{n=1}^{\infty}$ denote a sequence of self-adjoint operators on a Hilbert space \mathcal{H} and let F be a real-valued function on \mathbb{R} . We study here the question of whether $T_n = F(A_n)$ converging to a self-adjoint limit T ensures that A_n converges to a self-adjoint limit A . We are especially interested in the case for which F is not injective. In that case, we also require some additional condition on the convergence of $\{A_n\}$ in some weak sense, from which we conclude the convergence of $\{A_n\}$ in a stronger sense. More specifically, we introduce the notion of heat kernel regularization: this is obtained by regularizing the bilinear form A_n with a heat kernel $K_n = \exp(-T_n)$. The regularized operator is $K_n A_n K_n$. A norm convergent heat kernel regularization leads to a norm convergence of the resolvents of A_n .

Let us start with a simple, motivating example. Let $\mathcal{H} = \mathcal{K} \oplus \mathcal{K}$ be a direct sum of isomorphic subspaces, and let A_k have the off-diagonal form

$$A_k = \begin{pmatrix} O & Q_k^* \\ Q_k & O \end{pmatrix}$$

with respect to this decomposition. The function $F(t) = t^2$ gives rise to the positive operator

$$A_n^2 = \begin{pmatrix} Q_n^* Q_n & O \\ O & Q_n Q_n^* \end{pmatrix}.$$

It is clear that while A_n is a square root of A_n^2 , it is not a square root defined by the spectral theorem, which is diagonal with respect to $\mathcal{H} =$

* Supported in part by the National Science Foundation under Grant DMS/PHY 86-45122.

† Permanent address: Mathematics Department, ETH Zentrum, Zürich.

$\mathcal{X} \oplus \mathcal{X}$. The convergence $A_n^2 \rightarrow A^2$ does not in general yield $A_n \rightarrow A$, even if \mathcal{X} is finite-dimensional. This concrete example [2] led us to the more general question.

We are concerned here with norm-resolvent convergence. Let $R_n(z) = (A_n - z)^{-1}$ be the resolvent of A_n .

DEFINITION 1. The sequence of self-adjoint operators $\{A_n\}_{n=1}^\infty$ has a norm-resolvent limit A if for all z with $\text{Im } z \neq 0$, the operators $R_n(z)$ converge in norm to the resolvent $R(z)$ of a self-adjoint operator A .

Let us denote the norm-resolvent limit A of A_n by n.r. $\lim_{n \rightarrow \infty} A_n$. The two following results are well known.

PROPOSITION 2 [1]. *If $R_n(\pm i)$ converge in norm to operators R_\pm with densely defined inverses, then R_\pm is the resolvent of a self-adjoint operator A and n.r. $\lim A_n = A$.*

PROPOSITION 3 (Theorem VIII.20 of [3]). *Let g be a continuous function which vanishes at ∞ . Then n.r. $\lim A_n = A$ ensures that as $n \rightarrow \infty$,*

$$\|g(A_n) - g(A)\| = o(1).$$

In this note we look for a partial converse to Proposition 3. We study functions $g(t) = f(|t|)$, where f is well behaved:

DEFINITION 4. A continuous function $f: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is well behaved if f is monotonic and $\ln t \leq f(t)$ for t sufficiently large.

For any well-behaved f we define a function from self-adjoint operator A to "heat kernel" contraction semigroups K^β on \mathcal{H} by the map

$$A \mapsto K(A)^\beta = \exp(-\beta f(|A|)), \quad \beta \geq 0.$$

Let $K_n = K(A_n)$ be the semigroup for A_n . Since f increases monotonically to ∞ , f has a continuous inverse. Also $A_n K_n$ is a bounded operator, with $\|A_n K_n\|$ bounded uniformly in n . In particular, if $\delta A_{nm} = (A_n - A_m)^-$ is the closure of $A_n - A_m$, then $K_n \delta A_{nm} K_m$ is bounded for all n, m . Note that with our assumptions on f , the sequence $\{K_n A_n K_n\}$ is norm convergent if and only if $\|K_n \delta A_{nm} K_m\| = o(1)$ as $n, m \rightarrow \infty$.

DEFINITION 5. The sequence $\{A_n\}$ has a convergent heat kernel regularization with respect to f if

$$\|K_n \delta A_{nm} K_m\| = o(1),$$

as $n, m \rightarrow \infty$.

Our main result is the following:

THEOREM 6. *Let f be well behaved and let $\{A_n\}$ be a sequence of operators with a convergent heat kernel regularization with respect to f . Let $f(|A_n|)$ have a norm-resolvent limit. Then there exists a self-adjoint operator A such that*

$$T = \text{n.r. lim } f(|A_n|) = f(|A|),$$

and

$$\text{n.r. lim } A_n = A.$$

LEMMA 7. *Let $S_n = (A_n + i)^{-1}$. Let $\partial_n = \partial(|A_n|)$, where $\partial: [0, \infty) \rightarrow [1, \infty)$ is a monotonic, continuous function tending to ∞ , and satisfying $\partial(\lambda) = O(\lambda)$. Then $\partial_n S_n$ is bounded uniformly in n .*

Proof. By the spectral theorem, $|A_n|$ commutes with S_n , so the magnitude of the spectrum of $\partial_n S_n$ is

$$|(\lambda + i)^{-1}| \partial(|\lambda|) \tag{1}$$

for $\lambda \in \text{spectrum } A_n$. Since $\partial(|\lambda|) = O(|\lambda|)$, it follows that (1) is bounded. The bound is independent of n , and ensures $\|\partial_n S_n\| \leq \text{const}$.

Proof of the Theorem. The resolvent identity on the range of K_n is

$$\delta S = S_n - S_m = -S_m \delta A_{nm} S_n.$$

Let $E_n(\lambda)$ denote the spectral projection of $f(|A_n|)$ onto the subspace $f(|A_n|) \leq \lambda$. We study

$$\delta S = \delta S E_n + \delta S (I - E_n).$$

Using the lemma, as $\lambda \rightarrow \infty$, and letting ∂_n^{-1} denote the operator inverse of ∂_n (rather than the inverse function),

$$\begin{aligned} \|S_n(I - E_n)\| &= \|S_n \partial_n \partial_n^{-1}(I - E_n)\| = O(1) \|\partial_n^{-1}(I - E_n)\| \\ &= o(1), \end{aligned} \tag{2}$$

uniformly in n . Also

$$\begin{aligned} S_m(I - E_n) &= S_m \partial_m \partial_m^{-1}(I - E_n) \\ &= S_m \partial_m (\partial_m^{-1} - \partial_n^{-1})(I - E_n) + S_m \partial_m \partial_n^{-1}(I - E_n). \end{aligned}$$

Since $f(|A_n|)$ converges in the norm resolvent sense by hypothesis, it

follows that $h(|A_n|)$ is norm-convergent where h is any bounded function which vanishes at ∞ . In particular, $\|\partial_n^{-1} - \partial_m^{-1}\| = o(1)$ as $n, m \rightarrow \infty$. Thus

$$\begin{aligned} \|S_m(I - E_n)\| &= o(1) + O(1) \|\partial_n^{-1}(I - E_n)\| \\ &= o(1) + o(1). \end{aligned} \tag{3}$$

From (2) to (3) we infer that given $\varepsilon > 0$, we can choose λ_0, N such that for $\lambda > \lambda_0$ and $n, m > N$,

$$\|\delta S(I - E_n)\| \leq \varepsilon. \tag{4}$$

Likewise, taking the adjoint, and exchanging n, m ,

$$\|(I - E_m) \delta S\| \leq \varepsilon. \tag{5}$$

We have

$$\delta S E_n = E_m \delta S E_n + (I - E_m) \delta S E_n,$$

so from (4) to (5) we conclude

$$\|\delta S\| \leq 2\varepsilon + \|E_m \delta S E_n\|. \tag{6}$$

Now

$$\begin{aligned} E_m \delta S E_n &= -S_m E_m \delta A_{nm} E_n S_n \\ &= -S_m K_m^{-1} E_m K_m \delta A_{nm} K_n E_n K_n^{-1} S_n \end{aligned}$$

and

$$\|E_m \delta S E_n\| \leq e^{2\lambda} \|K_m \delta A_{nm} K_n\|.$$

Using the hypothesis, we choose n, m sufficiently large that

$$\|E_m \delta S E_n\| \leq \varepsilon, \tag{7}$$

By (6) and (7)

$$\|\delta S\| \leq 3\varepsilon,$$

and hence there exists S such that $\|S_n - S\| \rightarrow 0$. We can repeat the same estimates with S_n^* to conclude that $S_n^* \rightarrow S^*$.

We wish to conclude that S is the resolvent of a self-adjoint operator A . It is sufficient to show that S and S^* have densely defined inverses, and to use Proposition 2. Let $g(\cdot) = (f^{-1}(\cdot)^2 + I)^{-1}$. Since $f \nearrow \infty$ and f is invertible, we infer that $f^{-1} \nearrow \infty$. Thus $g(\cdot) \searrow 0$ at infinity. By

Proposition 3, $g(f(|A_n|)) = (A_n^2 + I)^{-1}$ has a limit $(f^{-1}(T)^2 + I)^{-1}$, where $T = \text{n.r. lim } f(|A_n|)$. But $S_n^* S_n = (A_n^2 + I)^{-1} = S_n S_n^*$. Thus

$$\lim_{n \rightarrow \infty} S_n^* S_n = \lim_{n \rightarrow \infty} S_n S_n^* = (f^{-1}(T)^2 + I)^{-1} = S^* S = S S^*$$

has a dense range and a densely defined inverse (with domain $D(f^{-1}(T)^2)$). Hence $\text{Range}(S) = \text{Domain}(S^{-1})$ and $\text{Domain}(S^*^{-1})$ are dense. This completes the proof of the theorem.

REFERENCES

1. J. GLIMM AND A. JAFFE, Singular perturbations of self-adjoint operators, *Comm. Pure Appl. Math.* **22** (1969), 401–414.
2. A. JAFFE, A. LESNIEWSKI, AND J. WEITSMANN, The two-dimensional $N = 2$, Wess–Zumino model on a cylinder, *Commun. Math. Phys.* **114** (1988), 147–165.
3. M. REED AND B. SIMON, "Methods of Modern Mathematical Physics I: Functional Analysis," Academic Press, New York, 1980.